

After placement of dental implant

Dr. Jason Hwang

Swelling is a normal occurrence after surgery. It will begin to subside after 48hrs. Use an ice pack.

If antibiotics are prescribed, continue to take them for the indicated length of time even if signs and symptoms of infection are gone.

Resume your normal dental routine after 24hrs, this will speed healing and help keep your mouth fresh and clean.

1. Bite on gauze for 45min after the appointment
2. Do not rinse vigorously
3. Do not suck on straws
4. Do not smoke
5. Do not drink alcohol
6. Do not brush teeth next to the extraction site for 72hrs
7. Limit vigorous exercise for the next 24hrs; this will increase blood pressure and may cause more bleeding
8. Take pain medications as prescribed
9. Drink lots of fluid and eat nutritious soft food on the day of the extraction; you can eat normally as soon as you are comfortable
10. Place an ice pack to the side of your face where surgery was performed

Call us if you have:

1. Heavy bleeding
2. Severe pain
3. Continued swelling for 2-3 days
4. Have a reaction to the medication

What do we mean by soft food diet?

Dr. Jason Hwang

For breakfast

Yogurt, eggs, soft crustless bread. Oatmeal or cold cereal will soak up milk and become mushy.

For Lunch and Dinner

Soup, mashed potatoes, pasta, macaroni and cheese, fish, tuna salad, egg salad.

Other soft foods

Applesauce, cottage cheese, Avocado, pudding, ice cream, muffins, milkshakes, fruit smoothies, jello.